

## Anti-SESN2 Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R012861

### Product Information

Application	WB, IHC-P/IF (Tissue-P), IF (Cell)/ICC, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	17M05M30
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant protein of human SESN2
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-SESN2 Rabbit mAb [17M05M30] is for research use only and not for use in diagnostic or therapeutic procedures.

### Protein Information

Synonyms	DKFZp761M0212; DKFZp761M02121; Hi95; Hypoxia induced gene 95; MGC11758; OTTHUMP0000003641; OTTMUSP00000010184; RP23 63D8.3; SES2; SESN 2; SESN2; SESN2_HUMAN; SEST2; Sestrin 2; Sestrin-2.
Calculated MW	Calculated MW: 55 kDa; Observed MW: 55 kDa
Uniprot ID	P58004
Gene ID	83667
Background	This gene encodes a member of the sestrin family of PA26-related proteins. The encoded protein may function in the regulation of cell growth and survival. This protein may be involved in cellular response to different stress conditions. [provided by RefSeq, Jul 2008]



Western Blot - Anti-SESN2 Rabbit mAb [17M05M30]

All lanes: R012861 at 1:1,000 dilution

Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2: Huh1 (Human hepatocarcinoma epithelial cell) whole cell lysates

Lane 3: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates

Lane 4: 293T (Human embryonic kidney cell) whole cell lysates

Lane 5: U87 (Human malignant glioblastoma epithelial cells) whole cell lysates

Lane 6: Raw264.7 (Mouse mononuclear macrophage leukemia cell) whole cell lysates

Lane 7: Mouse heart whole tissue lysates

Lane 8: Mouse liver whole tissue lysates

Lane 9: Mouse brain whole tissue lysates

Lane 10: PC-12 (Rat adrenal pheochromocytoma epithelial cell) whole cell lysates

Lane 11: Rat muscle whole tissue lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 55 kDa

Observed band size: 55 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-SESN2 Rabbit mAb [17M05M30]

Sample: Paraformaldehyde-fixed, paraffin embedded rat liver tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R012861 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-SESN2 Rabbit mAb [17M05M30]

Sample: Paraformaldehyde-fixed, paraffin embedded human colonic cancer tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R012861 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-SESN2 Rabbit mAb [17M05M30]

Sample: Paraformaldehyde-fixed, paraffin embedded mouse pancreas tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R012861 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



#### Immunofluorescence - Anti-SESN2 Rabbit mAb [17M05M30]

Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R012861 at 1:100 dilution and  $\alpha$ -tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).