

Anti-BAF57 Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R014050

Product Information

Application	IF (Cell)/ICC, ELISA, WB, IHC-P/IF (Tissue-P)
Reactivity	Human, Mouse, Rat
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	33K67P30
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human BAF57
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-BAF57 Rabbit mAb [33K67P30] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	BAF57, BRG1 associated factor 57, BRG1-associated factor 57, Chromatin remodeling complex BRG1 associated factor 57, FLJ35648, SMARCE 1, SMARCE1, SMCE1_HUMAN, SWI/SNF related matrix associated actin dependent regulator of chromatin e1, SWI/SNF related matrix associated actin dependent regulator of chromatin subfamily e member 1, SWI/SNF-related matrix-associated actin-dependent regulator of chromatin subfamily E member 1.
Calculated MW	Calculated MW: 47 kDa; Observed MW: 57 kDa
Uniprot ID	Q969G3
Gene ID	6605
Background	The protein encoded by this gene is part of the large ATP-dependent chromatin remodeling complex SWI/SNF, which is required for transcriptional activation of genes normally repressed by chromatin. The encoded protein, either alone or when in the SWI/SNF complex, can bind to 4-way junction DNA, which is thought to mimic the topology of DNA as it enters or exits the nucleosome. The protein contains a DNA-binding HMG domain, but disruption of this domain does not abolish the DNA-binding or nucleosome-displacement activities of the SWI/SNF complex. Unlike most of the SWI/SNF complex proteins, this protein has no yeast counterpart. [provided by RefSeq, Jul 2008]
Cellular Location	Nucleus.



Western Blot - Anti-BAF57 Rabbit mAb [33K67P30]

All lanes: R014050 at 1:1,000 dilution

Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates

Lane 3: HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates

Lane 4: A431 (Human epidermoid teratoma cell line) whole cell lysates

Lane 5: T24 (Human bladder cancer epithelial cell) whole cell lysates

Lane 6: Rat spleen whole tissue lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 47 kDa

Observed band size: 57 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-BAF57 Rabbit mAb [33K67P30]

Sample: Paraformaldehyde-fixed, paraffin embedded mouse kidney tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.


Primary antibody: R014050 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution
DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-BAF57 Rabbit mAb [33K67P30]

Sample: Paraformaldehyde-fixed, paraffin embedded human endometrial carcinoma tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.


Primary antibody: R014050 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution
DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunofluorescence - Anti-BAF57 Rabbit mAb [33K67P30]

Sample: HeLa cells

The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.5% Triton X-100 for 10 minutes and then blocked with 5% BSA in 0.1% PBS-Tween for 0.5 hours.

Primary antibodies: R014050 at 1:100 dilution and α-tubulin Mouse Monoclonal Antibody (Cat. No. LF209) at 1:100 dilution

Secondary antibodies: Goat anti-Rabbit (488) at 1:1,000 dilution (shown in green) and Goat anti-Mouse (555) at 1:1,000 dilution (shown in red)

Nuclei were stained with DAPI (shown in blue).