

Anti-TRADD Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody


Catalog # R016073

Product Information

Application	WB, IHC-P/IF (Tissue-P), ELISA
Reactivity	Human
Dilution	WB 1:1,000~1:2,000; IHC-P 1:100~1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	59Q72E33
Isotype	IgG
Label	Unconjugated
Immunogen	A synthesized peptide derived from human TRADD
Format	Affinity purified monoclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-TRADD Rabbit mAb [59Q72E33] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	Hs.89862; 9130005N23Rik; TRADD_HUMAN; TRADD; TNFR1-associated DEATH domain protein; TNFRSF1A-associated via death domain; TRADD_MOUSE.
Calculated MW	Calculated MW: 34 kDa; Observed MW: 34 kDa
Uniprot ID	Q15628
Gene ID	8717
Background	The protein encoded by this gene is a death domain containing adaptor molecule that interacts with TNFRSF1A/TNFR1 and mediates programmed cell death signaling and NF-kappaB activation. This protein binds adaptor protein TRAF2, reduces the recruitment of inhibitor-of-apoptosis proteins (IAPs) by TRAF2, and thus suppresses TRAF2 mediated apoptosis. This protein can also interact with receptor TNFRSF6/FAS and adaptor protein FADD/MORT1, and is involved in the Fas-induced cell death pathway. [provided by RefSeq, Jul 2008]
Cellular Location	Nucleus Cytoplasm Cytoplasm Cytoskeleton Shuttles between the cytoplasm and the nucleus.
Tissue Location	Found in all examined tissues.



Western Blot - Anti-TRADD Rabbit mAb [59Q72E33]

All lanes: R016073 at 1:1,000 dilution

Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates

Lane 2: HepG2 (Human hepatocarcinoma epithelial cell) whole cell lysates

Lane 3: HCT116 (Human colorectal carcinoma epithelial cell) whole cell lysates

Lane 4: 293T (Human embryonic kidney cell) whole cell lysates

Lane 5: MCF-7 (human breast adenocarcinoma epithelial cell) whole cell lysates

Lane 6: U87 (Human malignant glioblastoma epithelial cells) whole cell lysates


Lysates/proteins at 10 µg per lane.

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution

Predicted band size: 34 kDa

Observed band size: 34 kDa

Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-TRADD Rabbit mAb [59Q72E33]

Sample: Paraformaldehyde-fixed, paraffin embedded human breast cancer tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R016073 at 1:200 dilution


Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-TRADD Rabbit mAb [59Q72E33]

Sample: Paraformaldehyde-fixed, paraffin embedded human renal carcinoma tissue

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.

Primary antibody: R016073 at 1:200 dilution

Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution

DAB was used as the chromogen.

Counter stained with hematoxylin.

Positive/negative staining were presented.

Only the secondary antibody was used as the negative control.