Anti-AKT1/2/3 Rabbit mAb

Purified Recombinant Rabbit Monoclonal Antibody

Catalog # R011234


Product Information

Application	WB, IHC-P/IF (Tissue-P), ELISA
Reactivity	Human, Mouse
Dilution	WB 1:1,000~1:2,000; IHC-P 1:200; IF 1:100~1:200
Host	Rabbit
Clonality	Monoclonal
Clone No.	69K96M86
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant protein of human AKT1/2/3
Format	Affinity purified monoclonal antibody supplied in PBS with 0.01% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-AKT antibody [69K96M86] is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Synonyms	AKT1, PKB, RAC, RAC-alpha serine/threonine-protein kinase, Protein kinase B, PKB, Protein kinase B alpha, PKB alpha, Proto-oncogene c-Akt, RAC-PK-alpha.
Calculated MW	Calculated MW: 56 kDa; Observed MW: 55-60 kDa
Uniprot ID	P31749, P31751, Q9Y243
Gene ID	207/208/10000
RRID	AB_3752611
Background	Akt, also referred to as PKB or Rac, plays a critical role in controlling survival and apoptosis. This protein kinase is activated by insulin and various growth and survival factors to function in a wortmannin-sensitive pathway involving PI3 kinase. Akt is activated by phospholipid binding and activation loop phosphorylation at Thr308 by PDK1 and by phosphorylation within the carboxy terminus at Ser473.


Validation Images



Western Blot - Anti-AKT Rabbit mAb [69K96M86]
All lanes: R011234 at 1:1,000 dilution
Lane 1: HeLa (human cervix adenocarcinoma epithelial cell) whole cell lysates
Lane 2: MCF7 (human breast cancer epithelial cell) whole cell lysates
Lane 3: MSC (human mesenchymal stem cell) whole cell lysates
Lane 4: C2C12 (mouse myoblasts epithelial cell) whole cell lysates
Lysates/proteins at 10 µg per lane.
Secondary antibody: Goat Anti-Rabbit IgG(H+L), HRP Conjugated (Cat. No. LF102) at 1:5,000 dilution
Predicted band size: 56 kDa
Obsered band size: 55-60 kDa
Developed using the ECL technique (Cat. No. SQ201).



Immunohistochemistry - Anti-AKT Rabbit mAb [69K96M86]
Sample: Paraformaldehyde-fixed, paraffin embedded human hepatoma carcinoma tissue
Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.
Primary antibody: R011234 at 1:200 dilution
Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution
DAB was used as the chromogen.
Counter stained with hematoxylin.
Positive/negative staining were presented.
Only the secondary antibody was used as the negative control.



Immunohistochemistry - Anti-AKT Rabbit mAb [69K96M86]
Sample: Paraformaldehyde-fixed, paraffin embedded human cervical cancer tissue
Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0) for 30 mins.
Primary antibody: R011234 at 1:200 dilution
Secondary antibody: Goat Anti-Rabbit IgG (H+L), HRP conjugated at 1:1,000 dilution
DAB was used as the chromogen.
Counter stained with hematoxylin.
Positive/negative staining were presented.
Only the secondary antibody was used as the negative control.