Anti-p19ARF Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P105429


Product Information

Application	WB, ELISA
Reactivity	Human
Dilution	WB 1:500~1:5000
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	A synthetic peptide corresponding to a sequence within amino acids 1-100 of mouse p19ARF (NP_034007.1).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-p19ARF Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	Arf; p16; MTS1; Pctr1; p19ARF; p16INK4a; p19<ARF>; ARF-INK4a; INK4a-ARF; Ink4a/Arf; p16(INK4a).
Calculated MW	Calculated MW: 19 kDa; Observed MW: 17 kDa
Uniprot ID	Q64364
Gene ID	12578
Background	Enables MDM2/MDM4 family protein binding activity; enzyme inhibitor activity; and protein N-terminus binding activity. Involved in several processes, including negative regulation of lymphocyte proliferation; positive regulation of apoptotic process; and regulation of transcription, DNA-templated. Acts upstream of or within several processes, including negative regulation of cellular protein metabolic process; negative regulation of mammary gland epithelial cell proliferation; and positive regulation of apoptotic process involved in mammary gland involution. Located in cytoplasm and granular component. Part of protein-containing complex. Is expressed in several structures, including bone marrow; central nervous system; dorsal root ganglion; eye; and testis. Human ortholog(s) of this gene implicated in several diseases, including breast cancer (multiple); carcinoma (multiple); hematologic cancer (multiple); melanoma (multiple); and nervous system cancer (multiple). Orthologous to human CDKN2A (cyclin dependent kinase inhibitor 2A).

Validation Images



Western blot analysis of lysates from 293F cells, using p19ARF Rabbit pAb (P105429) at 1:1,000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Kit.
Exposure time: 15s.