[KO Validated] Anti-IKK gamma Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P101236


Product Information

Application	WB, IP, ELISA
Reactivity	Human, Mouse
Dilution	WB 1:500~1:2000; IP 0.5μg-4μg antibody for 200μg-400μg extracts of whole cells
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 1-419 of human IKKγ (NP_003630.1).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-IKK gamma Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Synonyms	IP; IP1; IP2; FIP3; IKKG; IPD2; NEMO; FIP-3; Fip3p; IMD33; SAIDX; AMCBX1; EDAID1; IKKAP1; ZC2HC9; IKK-gamma; IKKγ.
Calculated MW	Calculated MW: 48 kDa; Observed MW: 48 kDa
Uniprot ID	Q9Y6K9
Gene ID	8517
Background	This gene encodes the regulatory subunit of the inhibitor of kappaB kinase (IKK) complex, which activates NF-kappaB resulting in activation of genes involved in inflammation, immunity, cell survival, and other pathways. Mutations in this gene result in incontinentia pigmenti, hypohidrotic ectodermal dysplasia, and several other types of immunodeficiencies. A pseudogene highly similar to this locus is located in an adjacent region of the X chromosome.


Validation Images



Western blot analysis of lysates from wild type (WT) and IKKγ knockout (KO) 293T cells, using [KO Validated] IKKγ Rabbit pAb (P101236) at 1:1,000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Kit.
Exposure time: 5s.



Western blot analysis of various lysates using [KO Validated] IKKγ Rabbit pAb (P101236) at 1:1,000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.



Immunoprecipitation analysis of 200 µg extracts of A549 cells using 3 µg IKKγ antibody (P101236). Western blot was performed from the immunoprecipitate using IKKγ antibody (P101236) at a dilution of 1:1000.