Anti-Filamin B Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P105479


Product Information

Application	WB, IHC-P, IF (Cell)/ICC, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:500~1:1000; IHC-P 1:50~1:100; IF 1:20~1:100
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 1686-1785 of human FLNB (NP_001157789.1).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Filamin B Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Synonyms	AOI; FH1; SCT; TAP; LRS1; TABP; FLN-B; FLN1L; ABP-278; ABP-280; FLNB.
Calculated MW	Calculated MW: 278 kDa; Observed MW: 298 kDa
Uniprot ID	O75369
Gene ID	2317
Background	This gene encodes a member of the filamin family. The encoded protein interacts with glycoprotein Ib alpha as part of the process to repair vascular injuries. The platelet glycoprotein Ib complex includes glycoprotein Ib alpha, and it binds the actin cytoskeleton. Mutations in this gene have been found in several conditions: atelosteogenesis type 1 and type 3; boomerang dysplasia; autosomal dominant Larsen syndrome; and spondylocarpotarsal synostosis syndrome. Multiple alternatively spliced transcript variants that encode different protein isoforms have been described for this gene.


Validation Images




Immunohistochemistry analysis of paraffin-embedded Rat heart using FLNB Rabbit pAb (P105479) at dilution of 1:100 (40× lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human liver damage using FLNB Rabbit pAb (P105479) at dilution of 1:100 (40× lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.




Immunohistochemistry analysis of paraffin-embedded Human prostate using FLNB Rabbit pAb (P105479) at dilution of 1:100 (40× lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.




Immunofluorescence analysis of HeLa cells using FLNB Rabbit pAb (P105479) at dilution of 1:100 (40× lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of HUVEC cells using FLNB Rabbit pAb (P105479) at dilution of 1:100 (40× lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 cells using FLNB Rabbit pAb (P105479) at dilution of 1:100 (40× lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of PC-12 cells using FLNB Rabbit pAb (P105479) at dilution of 1:100 (40× lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates using FLNB Rabbit pAb (P105479) at 1:1,000 dilution._Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution._Lysates/proteins: 25µg per lane._Blocking buffer: 3% nonfat dry milk in TBST._Detection: ECL Kit._Exposure time: 10s.