Anti-Rac1/2/3 Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P108103


Product Information

Application	WB, IHC-P, IF (Cell)/ICC, ELISA
Reactivity	Human, Mouse, Rat
Dilution	WB 1:500~1:1000; IHC-P 1:50~1:200; IF 1:50~1:200
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	Recombinant fusion protein containing a sequence corresponding to amino acids 1-192 of human Rac1/2/3 (NP_008839.2).
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Rac1/2/3 Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Synonyms	MIG5; MRD48; Rac-1; TC-25; p21-Rac1; Rac1/2/3.
Calculated MW	Calculated MW: 21 kDa; Observed MW: 21 kDa
Uniprot ID	P63000
Gene ID	5879
Background	The protein encoded by this gene is a GTPase which belongs to the RAS superfamily of small GTP-binding proteins. Members of this superfamily appear to regulate a diverse array of cellular events, including the control of cell growth, cytoskeletal reorganization, and the activation of protein kinases. Two transcript variants encoding different isoforms have been found for this gene.


Validation Images




Immunohistochemistry analysis of paraffin-embedded Human gastric using Rac1/2/3 Rabbit pAb (P108103) at dilution of 1:100 (40× lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.




Immunofluorescence analysis of C6 cells using Rac1/2/3 Rabbit pAb (P108103) at dilution of 1:100 (40× lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of NIH-3T3 cells using Rac1/2/3 Rabbit pAb (P108103) at dilution of 1:100 (40× lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of C6 cells using Rac1/2/3 Rabbit pAb (P108103) at dilution of 1:100 (40× lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 cells using Rac1/2/3 Rabbit pAb (P108103) at dilution of 1:100 (40× lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates using Rac1/2/3 Rabbit pAb (P108103) at 1:1,000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Kit.
Exposure time: 30s.



Western blot analysis of lysates from A-431 cells, using Rac1/2/3 Rabbit pAb (P108103) at 1:1,000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Kit.
Exposure time: 90s.