Anti-Phospho-PKC theta (Ser676) Rabbit pAb

Purified Rabbit Polyclonal Antibody

Catalog # P108802


Product Information

Application	WB, ELISA
Reactivity	Human
Dilution	WB 1:500~1:1000
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Label	Unconjugated
Immunogen	A phospho specific peptide corresponding to residues surrounding S676 of human PRKCQ
Format	Affinity purified polyclonal antibody supplied in PBS with 0.02% sodium azide and 50% glycerol, pH 7.3.
Storage	Shipped on wet ice. Store at -20°C. Stable for 24 months from date of receipt. Aliquoting is unnecessary for -20°C storage.
Precautions	Anti-Phospho-PKC theta (Ser676) Rabbit pAb is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Synonyms	PRKCT; PRKCQ; nPKC-theta; Phospho-PRKCQ-S676.
Calculated MW	Calculated MW: 82 kDa; Observed MW: 78 kDa
Uniprot ID	Q04759
Gene ID	5588
Background	Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role. The protein encoded by this gene is one of the PKC family members. It is a calcium-independent and phospholipid-dependent protein kinase. This kinase is important for T-cell activation. It is required for the activation of the transcription factors NF-kappaB and AP-1, and may link the T cell receptor (TCR) signaling complex to the activation of the transcription factors.

Validation Images



Western blot analysis of lysates from Jurkat cells, using Phospho-PRKCQ-S676 Rabbit pAb (P108802) at 1:1,000 dilution. Jurkat cells were treated by PMA/TPA (200 nM) at 37°C for 30 minutes after serum-starvation overnight.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (LF102) at 1:10,000 dilution.
Lysates/proteins: 25μg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Kit.
Exposure time: 60s.